



Expression of *Cyp1a* and *p53* Induced by Water-Soluble Fractions of Crude Oil in Juveniles of *Clarias gariepinus* (Burchell, 1822).

¹Gbuvboro, A. E, ^{2*}Eriegha, O. J and ²Eyo, V. O

¹Department of Environmental Management and Pollution

²Department of Fisheries and Aquaculture

Nigeria Maritime University, Okerenkoko, Nigeria.

Article Information

Article # 10002
Received: 9th Augt 2023
1st Revision: 1st Jan. 2023
2nd Revision: 4th Feb. 2024
Acceptance: 9th Feb. 2024
Available online:
3rd March 2024

Key Words

Oil pollution, Gene expression, *Clarias gariepinus* Cytochromes P4501A, p53.

Abstract

Oil spills, like most other sources of pollutants, have been shown to disrupt biochemical pathways and cause the death of organisms. In this study, the relative expression of *Cyp1a* and *p53* genes in juveniles of *Clarias gariepinus* was determined after exposure to sub-lethal concentrations of water-soluble fractions of crude oil. Fish (5.1 ± 0.3 g) were exposed to 4 sublethal concentrations (30, 45, 60, and 75% of the LC₅₀ corresponding to 67, 101, 135, and 169 mg/L respectively) of the oil and a control. After 90 days of exposure, fish were sacrificed, tissue collected, and used for gene expression analysis. Total RNA was extracted using a modified CTAB extraction protocol. Gene quantification and amplification were carried out using standard methods. The *B-actin* gene was used as an internal control. The results revealed that the relative expression of *Cyp1a* mRNA increased in a dose-dependent manner, indicating the upregulation of *Cyp1a*. In contrast to *Cyp1a*, crude oil modulated the expression of *p53*, but the expression pattern did not vary in a dose-dependent manner. The expression of the *Cyp1a* gene can therefore be used as an important factor for assessing the toxicity of crude oil, especially at sub-lethal concentrations in fish species.

*Corresponding Author: Eriegha, O. J.; oj.eriegha@yahoo.com

Introduction

Oil pollution is the introduction of liquid petroleum hydrocarbons into the environment, primarily from human activities, and is a major environmental concern (Adesanmi *et al.*, 2021). In Nigeria, the majority of the oil pollution occurs as a result of shipping and terrestrial traffic accidents, improper disposal of drilling muds, depot leakage, tank washing and oil ballast discharges, failure or rupture in oil pipelines, and in the process of oil theft or artisanal refining (bunkering) of crude oil (Ayuba, 2012; Kadafa *et al.*, 2012). This can have cascading effects on the entire ecosystem and reduction of biodiversity (Eriegha and Sam, 2020). Studies have shown that oil pollution can lead to the death of fish, birds, and marine mammals due to direct toxicity or smothering effects. It can also disrupt important biochemical pathways, such as oxidative stress and detoxification mechanisms, leading to cellular damage and impaired physiological functions (Eriegha *et al.*, 2019; Eriegha, 2023; Eriegha *et al.*, 2023). Additionally, exposure to oil can cause DNA damage and chromosomal abnormalities, affecting the development and growth of organisms, including

embryos and larvae (Santos *et al.*, 2016; Ardeshir *et al.*, 2018). Oil spills can also contaminate water sources, making them unsafe for human use and damaging the fishing and tourism industries, which are often important sources of income for local communities (Andrews *et al.*, 2021).

Recent advances in bioassay have expanded beyond traditional endpoints (such as enzymatic assays and protein quantification) to include gene expression studies. While enzymatic assays and protein quantification provide valuable information on the biochemical effects of toxic substances, gene expression studies offer insights into the molecular mechanisms underlying toxicity. Several studies have demonstrated the effectiveness of gene expression studies in toxicology using RT-qPCR (Piña *et al.*, 2007; Chivittz *et al.*, 2016; Santos *et al.*, 2018). By evaluating mRNA levels using techniques like RT-qPCR, researchers can identify specific genes that are upregulated or downregulated in response to exposure. This information can help identify potential biomarkers for toxicity and provide a deeper

understanding of the biological processes affected by toxic substances. The RT-QPCR is an effective tool for gene expression studies due to the accuracy, sensitivity and speed of results (Derveaux *et al.*, 2010).

The expression of genes such as Cytochromes P4501A (CYP1A) and p53, which plays a major role in the biotransformation of a variety of endogenous substances such as steroids, lipids, and vitamins, as well as environmental toxicants, is a veritable instrument for establishing the dynamics of the environment on cellular expression patterns (Kassahn *et al.*, 2007; Ardeshir *et al.*, 2018). The induction of *Cyp1a* by the various xenobiotics has led to its use as a biomarker in fish for assessing the level of contamination in an aquatic environment (Cousinou *et al.*, 2000; Hassanin *et al.*, 2009; Uno *et al.*, 2012). The p53 gene, also known as tumour protein, encodes a protein that regulates the cell cycle and so works as a tumor suppressor. Tumor suppression is a critical process in multicellular organisms to prevent uncontrolled cell proliferation and the development of cancer (Vogelstein *et al.*, 2000). As a transcription factor, p53 regulates the transcription of various genes involved in critical cellular processes. For example, it activates the transcription of *gadd45*, *p21*, and *cyclinG1*, which are genes responsible for cell cycle arrest. By upregulating these genes, p53 halts the progression of the cell cycle, allowing time for DNA repair or apoptosis if the damage is too severe (Speidel, 2015). Additionally, p53 also regulates the transcription of genes involved in DNA repair, such as *DDB2* and *XPC*, ensuring the proper maintenance of the genome. Furthermore, p53 can induce apoptosis by activating the transcription of genes like *bax* and *bcl-2*, which play crucial roles in programmed cell death. Understanding the transcriptional regulatory functions of p53 provides insights into how it controls key cellular processes to maintain cellular integrity (Liebermann and Hoffman, 2008).

Fish are good indicators of water contamination in aquatic systems for several reasons. Firstly, they occupy different trophic levels, and therefore represent different positions in the food chain. This is important because contaminants can accumulate and biomagnify as they move up the food chain, making higher trophic level organisms more likely to show signs of contamination (Odiete, 2003). Additionally, fish are more sensitive to many toxicants compared to invertebrates, making them more likely to exhibit

visible effects of contamination (Raimondo *et al.*, 2007). Lastly, fish have the ability to accumulate contaminants in their tissues over time, providing a record of exposure to specific pollutants (Hermoso *et al.*, 2010).

Clarias gariepinus is widely regarded as one of the most significant tropical catfish species for aquaculture, and is believed to have met all the attributes outlined for an ideal biosensor of environmental pollutants. These attributes may include the species' sensitivity to pollutants, its ability to accumulate contaminants in its tissues, and its responsiveness to changes in environmental conditions. This species has already been used in laboratory investigations and is a suitable organism for assessing the impact of xenobiotics (Olaifa, 2012; Obemeata *et al.*, 2012). Several studies have shown that sublethal exposure to contaminants can lead to physiological and behavioral changes in fish. However, much is still unclear regarding the consequences of low-level exposure on organisms, particularly at the molecular level. Sublethal effects are essential for predicting habitat and fish population impacts accurately, since many more organisms are likely to be exposed to sublethal concentrations of contaminants than to lethal concentrations as a spill disperses, and many may experience chronic low levels of one or more contaminants for a long time. Hence, this study aimed to determine the relative expression of *Cyp1a* and p53 induction in juveniles of *C. gariepinus* after exposure to water soluble fractions of crude oil. The findings of this study will provide valuable insights into the potential impact of crude oil exposure on the genetic expression.

Materials and Methods

Water-Soluble Fractions of Crude oil

Crude oil was obtained from the Afiesere oil field, was transported to the Department of Chemistry of the University of Ibadan where water soluble fraction (WSF) of the oil was prepared using Anderson *et al.* (1974) with little modifications. Briefly, 1 ml oil was mixed with 9 ml of freshwater. The mixture was slowly stirred with a magnetic stirrer for a period of 20 h at room temperature. The bottle was capped with aluminum foil to minimize evaporation of the more volatile oil hydrocarbons. After mixing, the oil and water phases were allowed to separate for to 6 h before the water phase was siphoned off, and utilized in experiments. Different concentrations of the WSF were prepared by diluting the stock WSF with water.

Experimental Fish

Healthy juveniles of *Clarias gariepinus* (average weight = 5.1 ± 0.3 g) with a good physical appearance were purchased from Elivital fish farm, Isiokolo and transported to the laboratory of the Department of Fisheries and Aquaculture of the Nigeria Maritime University, Okerenkoko in oxygen bags. Health status of the fish was ascertained based on the absence of lesions and other morphological diagnostic symptoms as suggested by Wedemeyer (1997). The experimental fish were first kept in the laboratory for 2 weeks in natural photo-regime conditions in 30 L experimental tanks containing tap water. The juveniles of *C. gariepinus* were randomly allocated to triplicate groups with 20 fish per tank. To prevent the fish from escaping, the experimental tanks were fitted with a lid constructed of fine polyethylene gauze screen with a mesh size of 1mm. This type of lid also allowed for proper ventilation within the tanks and ensured that no unwanted materials entered the tanks, and thereby maintaining a clean and controlled environment for the experiments. They were fed with a commercial feed, Coppens® *ad libitum* two times a day during the acclimatization period.

Preliminary Bioassay

A preliminary toxicity test was conducted in a static exposure system to determine the lethal concentration 50 (LC₅₀). Specimens of *C. gariepinus* (n = 20 per tank) were randomly allocated in 15 litres of the test solution. The following concentrations of WSF of crude oil were used for the definitive (acute) test: 0, 100, 180, 320, and 576 mg/L. All treatments and control were conducted in triplicate. The set up was not aerated and fish mortality was documented every 24 h for 96 h. Fish were considered dead when they remained immobile on the bottom of the tank or suspended in the water column even after mechanical stimuli. Dead fish were removed immediately in order to avoid contamination of experimental setup. Mortalities recorded were expressed as percentages of the test populations and the median lethal concentration (LC₅₀) values was calculated by using regression equation method of probit analysis (Sakuma, 1998).

Exposure Assessment

The juveniles of *C. gariepinus* were treated with four different sub-lethal doses (30, 45, 60 and 75% of the LC₅₀ (267 mg/L) corresponding to 67, 101, 135, and 169mg/l respectively) of WSF of crude oil and a control group. The test was carried out utilizing a semi-static renewal method in which the exposure medium was refreshed every three days to preserve the toxicant's strength while reducing the level of ammonia. The experimental fish were fed with a commercial feed, Coppens® *ad libitum* two times a day. After 90 days of treatment, fish were sacrificed, tissue collected and transported to the Biorepository and Clinical Virology Laboratory of the University College, Ibadan, Nigeria for gene expression analysis.

Analysis of *Cyp1a* and *p53* Gene Expression

RNA was isolated using an RNA extraction kit (YT9065, Yekta, Iran) following the manufacturer's instructions with little modification based on Santos *et al.* (2018). Briefly, fish tissue was thawed and approximately 100 mg of fish tissue was ground and mixed in nuclease-free tubes with 350 µL of RB buffer and 3.5 µL of beta-mercaptoethanol. The mixture was then passed through a filter column and centrifuged at 18,000g for 2 minutes. The supernatant was combined with 70% ethanol, passed through an RB mini-column, and centrifuged for 1 minute. To remove DNA contamination, 250 µL of washing buffer 1 was added to the column and centrifuge for 1 minute. The column membrane was then subjected to 60 µL of RNase-free DNase I solution (Roche) for 15 minutes. After cleaning and drying the column twice, ddH₂O was added to it and centrifuged for one minute. To control the quality of extracted RNA, 2 µL of loading buffer and 5 µL of the RNA were loaded on 1% electrophoresis gel and electrophoresed for 30 min (85 V). A Nanodrop spectrophotometer (Thermo Scientific®) was used to measure the RNA concentration and purity. The absorbance of the RNA sample was measured at wavelengths of 230 nm, 260 nm, and 280 nm. cDNA synthesis was achieved using a commercially available kit (Thermo Scientific®). RiboLock RNase Inhibitor (1 µL) was applied after mixing and centrifuging random hexamer primer and RNA at 65 °C for 5 minutes (on ice). Then, RevertAid RT (1 µL), Reaction Buffer (4 µL), and dNTP (2 µL) were added. The reaction took place in a thermocycler (Biorad®) for 5 minutes at 25°C, 1 hour at 42°C, 5 minutes at 70°C, and an indefinite time at 20°C.

To design specific qPCR primers for the specific *Clarias gariepinus* genes, mRNA reference sequences of individual genes were obtained from the genbank database on the NCBI website (National Center for Biotechnology Information) site was then accessed and sequence pasted in the sequence entry box and multiple intercalating dye PCR primers were generated. It is very necessary to ensure that the primers will have a perfect match, this will enhance primer annealing during PCR. To do this, primers must anneal to regions where the sequences are conserved. Each primer pair was then checked for specificity to be sensitive to only the genes of interest to which it was designed to detect and also the ability to cut across all aligned genes then the best primer was selected and synthesized at Inqaba Biotech Laboratory in South Africa. The forward and reverse sequences of designed primers are presented in Table 1.

For the treatment of RNA, 20 ng total RNA was treated with NEB DNase 1 (M0303) to eliminate extracted DNA. A mixture of 2 µl of 10ng/ µl RNA, 10 µl DNase I Reaction Buffer (10X), 1 µl DNase I (RNase-free) and up to 100 µl with Nuclease-free H₂O. After 10 minutes of incubation at 37°C, 1 µl of 0.5 M EDTA (to a final concentration of 5 mM) was added. The mixture was then heat-inactivated at 75°C for 10 minutes before being stored at -20°C until use. Gene quantification was done following the Luna® Universal qPCR Master Mix Protocol (M3003). This was also used to detect the presence of mRNA genes in the extracted RNA. Expression of the B-actin gene was used as an internal control. A mix of 10 µl Luna Universal qPCR Master Mix, 0.5 µl Forward primer (10 µM), 0.5 µl Reverse primer (10 µM) and 0.06 Reverse Transcriptase (Promega) made up to 18 µl with Nuclease-free Water to which 2 µl of the treated RNA Template was added. The mixture was run with the profile initial denaturation at 95°C for 60 seconds followed by 40-45 of Denaturation at 95°C 15 seconds extension and plate reading at 60°C for 30 seconds and thereafter a termination at 72°C for 10 minutes. Amplification was conducted using the cfx96tm real-time system from bio-rad following the manufacturer's manual. The relative expression of *Cyp1a* and *p53* genes was computed using the comparative CT ($2^{-\Delta\Delta CT}$) method of Pfaffl *et al.* (2002).

Results and Discussion

The result of the quality assessment from the Nanodrop spectrophotometer is presented in Table 2. The

quantity assessment of RNA from the Nanodrop spectrophotometer showed that A260/280 ranged from 1.40 – 1.68 and A260/230 ranged from 1.62 – 1.92, implying suitable purity of the RNA.

The result of relative *Cyp1a* and *p53* mRNA expression in the tissue of *C. gariepinus* is presented in Figure 1. The result revealed that the relative expression of *Cyp1a* increased (upregulation) in a dose-dependent manner. The expression level increased as the concentration of the water-soluble fractions of crude oil increased. The control group had the least (3.39) while the highest relative expression (21.37) was observed in the group containing the highest concentrations of the toxicant. The upregulation of *Cyp1a* is crucial for xenobiotic metabolism and an indication of the activation of the aryl hydrocarbon receptor (AHR) pathway. This is because, polycyclic aromatic hydrocarbons (considered the most toxic component of crude oil) metabolism is mediated by the AHR, a ligand-binding transcription factor with a high affinity for PAH and polychlorinated biphenyl ligands (Billiard *et al.* 2002). Many genes in the AHR gene battery such as *Cyp1a* are involved in xenobiotic clearance (Dalton *et al.*, 2002; Jones, 2019). Induction of *Cyp1a* transcription is such a ubiquitous action of AHR activation that its upregulation is considered a classic biomarker of contaminant exposure. *Cyp1a* is a well-known biomarker of exposure to PAH (Lee and Anderson, 2005). Hook *et al.* (2010) have reported the upregulation of *Cyp1a* by crude oil, heavy fuel oil, and gas oil, but not by kerosene. Because kerosene does not contain substantial concentrations of high molecular weight PAH. Sustained activation of the *Cyp1a* gene was predictive of persistent exposure to sublethal concentrations of crude oil components and negative population-level impacts in fish following the Exxon Valdez oil spill (Whitehead *et al.*, 2012).

The findings of this study were in agreement with Sorhus *et al.* (2021) who reported a stable expression of *Cyp1a* in embryonic development in *Melanogrammus aeglefinus* after exposure to crude oil toxicity. They however reported a dynamic temporal expression of *Cyp1b*, *Cyp1c*, and *Cyp1d* indicating that these enzymes are actively involved in the metabolism of xenobiotics. Specifically, they are responsible for the biotransformation of these foreign compounds, converting them into more soluble forms that can be easily excreted. Upregulation in *Cyp1a* expression has also been reported in southern flounder (*Paralichthys*

lethostigma) after exposure to crude oil. Elevated expression of *Cyp1a* is indicative of hydrocarbon metabolism and is a commonly used biomarker for exposure to oil (Bilbao *et al.*, 2010).

Crude oil exposure has been shown to trigger gene pathways involved in biotransformation such as *cyp1a* and *cyp1c1* mRNA expression, lipid metabolism, stress response, and ion regulation in early life stages of Atlantic haddock (Sørhus *et al.* 2017). Aranguren-Abadía *et al.* (2020) have also reported the expression of *Cyp1a* in the cardiac tissues, skin, and the developing liver of *Gadus morhua* at 8 and 10 days' post fertilization. *Cyp1a* also been shown to be protective and necessary for detoxification (Wincent *et al.*, 2015) and may also result in the production of some reactive toxic metabolites. Nebert *et al.* (2004) have also noted other enzymes that can produce metabolites and increase the toxicity. Myocardial *Cyp1a* induction was observed in *Clupea pallasii* exposed to crude oil (Incardona *et al.*, 2019).

The results of this study showed that water-soluble fractions of crude oil modulated the expression of p53 but the expression pattern did not vary in a dose-dependent manner (Figure 1). The lowest relative expression level (1.88) of p53 was observed in the group containing no crude oil while the highest relative expression (8.20) was observed in the group containing 45% of the LC₅₀. The influence of crude oil on the expression of the p53 gene can vary depending on the specific circumstances and exposure conditions. Studies have revealed that exposure to crude oil or its components can affect the expression of genes involved in various cellular processes, including those related to DNA repair and cell cycle regulation, such as the p53 gene. However, the exact mechanisms and outcomes of crude oil exposure on p53 gene expression can be complex and context-dependent. Conflicting results showing both upregulation and downregulation (decreased expression) of p53 in response to exposure to xenobiotics have been reported by many authors (Liu *et al.*, 2011). The specific composition of the crude oil, the concentration and duration of exposure, and the cell or tissue type being studied can all influence the observed effects on p53 gene expression. Mitra *et al.* (2018) have reported the down-regulation of p53 in *Rita rita* from a polluted riverine environment. Similar down-regulation of p53 has also been reported in Japanese medaka after 2 days of exposure to nonylphenol and bisphenol (Min *et al.*,

2003). Similar findings have also been reported in adult hermaphroditic fish; *Kryptolebias marmoratus* on exposure to endocrine-disrupting chemicals (Lee *et al.*, 2008). On the other hand, Willams and Hubberstey (2014) demonstrated upregulation of p53 in brown bullheads from contaminated regions, compared to clean sites. Similarly, up-regulation of p53 has been reported in different tissues of *Mytilus edulis* in response to environmental genotoxicants (Di *et al.*, 2011). Studies have suggested that p53 might be activated in fish due to genotoxic stress (Rau-Embry *et al.*, 2006).

Conclusion

Crude oil is a compound mixture of hydrocarbons that can have a significant impact on gene expression in fish. Exposure to crude oil can lead to the upregulation of genes involved in detoxification, stress response, and cell death. The effects of crude oil exposure on gene expression can be complex and long-lasting. These effects can lead to a variety of negative consequences, including reduced growth, impaired reproduction, and increased susceptibility to disease. A comparative assessment of the target genes revealed that the relative expression of *Cyp1a* was upregulated in a dose-dependent manner. Despite the modulation of expression in p53, there was no definite pattern, as expression did not vary in a dose-dependent manner. The study therefore recommends the use of *Cyp1a* as a suitable gene in the assessment of crude oil toxicity. Further research is needed to better understand the mechanisms of the regulation of the p53 gene.

Acknowledgements

We sincerely acknowledge the Tertiary Education Trust Fund (TETFUND) for providing the Institutional Based Research (IBR) Grant through the Nigeria Maritime University, to fund this research. Special thanks to the Vice Chancellor of Nigeria Maritime University, Prof. E. M. Adigio, the Director of Academic Planning; Prof. J.T. Tuaweri and the Dean of the Faculty of Environmental Management, Prof. P. A. Bariweni for the encouragement and support.

References

- Adesanmi, A.J., Okedere, O.B., Sonibare, J. A., Elehinafe, F. B., and Fakinle, B. S. (2021). Atmospheric particulate fractions from Nigerian crude oil spillage. *Environmental Challenges*. 5: 1 – 5.
- Anderson, J. W, Neef, J. M., Cox, B. A., Tatem, H. E. and Hightower, G. M. (1974). Characteristics of

dispersions and water soluble extracts of crude and refined oils and their toxicity to estuarine crustaceans and fish. *Marine Biology*. 27: 75 – 88.

Andrews, N., Bennett, N. J., Le Billon, P., Green, S. J., Cisneros-Montemayor, A. M., Amongin, S. and Sumaila, U. R. (2021). Oil, fisheries and coastal communities: A review of impacts on the environment, livelihoods, space and governance. *Energy Research & Social Science*. 75, 102009.

Aranguren-Abadía, L., Donald, C. E., Eilertsen, M., Gharbi, N., Tronci, V., Sørhus, E., Mayer, P., Nilsen, T. O., Meier, S., Goksøyr, A. and Karlsen, O. A. (2020). Expression and localization of the aryl hydrocarbon receptors and cytochrome P450 1A during early development of Atlantic cod (*Gadus morhua*). *Aquat. Toxicol.* 226, 105558.

Ardeshir, R. A., Zolgharnein, H., Movahedinia, A., Salamat, N., and Zabihi, E. (2018). CYP1A gene expression as a basic factor for fipronil toxicity in Caspian kutum fish. *Toxicology Reports*. 5: 113-124.

Ayuba, A. K. (2012). Environmental Impacts of Oil Exploration and Exploitation in the Niger Delta of Nigeria. *Global Journal of Science Frontier Research Environment and Earth Sciences*. 12(3): 18-28.

Bilbao, E., Raingeard, D., de Cerio, O. D., Ortiz-Zarragoitia M., Ruiz, P. and Izagirre, U. (2010). ZEffects of exposure to Prestige-like heavy fuel oil and to perfluorooctane sulfonate on conventional biomarkers and target gene transcription in the thicklip grey mullet *Chelon labrosus*. *Aquat Toxicol.* 98(3): 282–96.

Billiard, S. M., Hahn, M. E., Franks, D. G., Peterson, R. E., Bols, N. C. and Hodson, P. V. (2002). Binding of polycyclic aromatic hydrocarbons (PAHs) to teleost aryl hydrocarbon receptors (AHRs). *Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology*. 133(1): 55 – 68.

Chivittz, C. C., Pinto, D. P., Ferreira, R. S., Sopezki, M. D. S., Fillmann, G., and Zanette, J. (2016). Responses of the CYP1A biomarker in *Jenynsia multidentata* and *Phalloceros caudimaculatus* and evaluation of a CYP1A refractory phenotype. *Chemosphere*. 144: 925 - 931.

Cousinou, M., Nilsen, B., Lopez-Barea, J. and Dorado, G. (2000). New methods to use fish cytochrome P4501A to assess marine organic

pollutants. *Science of the total environment*. 247(2-3): 213 - 225.

Dalton, T. P., A. Puga, and H. G. Shertzer. (2002). Induction of cellular oxidative stress by aryl hydrocarbon receptor activation. *Chemico-Biological Interactions*. 141(1–2): 77 – 95.

Derveaux, S., Vandesompele, J. and Hellemsans, J. (2010). How to do successful gene expression analysis using real-time PCR. *Methods*. 50(4): 227 – 230.

Di, Y., Schroeder, D.C., Highfield, A., Readman, J.W. and Jha, A.N. (2011). Tissue-specific expression of p53 and ras genes in response to environmental genotoxicant benzo(α) pyrene in marine mussels. *Environ. Sci. Technol.* 45 (20): 8974 – 8981.

Eriegha O. J. and Sam K. (2020). Characterization of crude oil impacts and loss of livelihood in the Niger Delta, Nigeria: A fisheries perspective. *International Journal of Maritime and Interdisciplinary Research*. 1(1): 255 – 273.

Eriegha, O. J., Omitoyin, B. O. and Ajani, E. K. (2019). Water soluble fractions of crude oil deteriorates water quality parameters and alters histopathological components of juvenile *Clarias gariepinus*. *Animal Research International*. 16(2): 3308 – 3318.

Eriegha, O. J. (2023). Haematological and serum biochemical changes in *Clarias gariepinus* exposed to sub-lethal concentrations of water-soluble fractions of crude oil. *Nigerian Journal of Science and Environment*. 21(1): 261 – 270.

Eriegha, O. J., Eyo, V. O. and Adeleke, I. A. (2023). Influence of crude oil exposure and recovery on growth performance in *Oreochromis niloticus*. *Proceedings of the 1st international conference of the school of Agriculture, Food and Natural Resources*, Olusegun Agagu University of Science and Technology, Okitipupa, Ondo State. 13 – 16th November, 2023.

Hassanin, A., Ropiquet, A., Couloux, A., and Cruaud, C. (2009). Evolution of the mitochondrial genome in mammals living at high altitude: new insights from a study of the tribe Caprini (Bovidae, Antilopinae). *Journal of molecular evolution*. 68: 293 – 310.

Hermoso, V., Clavero, M., Blanco-Garrido, F. and Prenda, J. (2010). Assessing the ecological status in species-poor systems: A fish-based index for Mediterranean Rivers (Guadiana River, SW Spain). - *Ecological Indicators*. 10(6): 1152 – 1161.

- Holth, T. F., Nourizadeh-Lillabadi, R., Blæsbjerg, M., Grung, M., Holbech, H., Petersen, G. I. and Hylland, K. (2008). Differential gene expression and biomarkers in zebrafish (*Danio rerio*) following exposure to produced water components. *Aquatic toxicology*. 90(4): 277-291.
- Hook, S. E., Lampi, M. A., Febbo, E. J., Ward, J. A., and Parkerton, T. F. (2010). Hepatic gene expression in rainbow trout (*Oncorhynchus mykiss*) exposed to different hydrocarbon mixtures. *Environmental Toxicology and Chemistry*. 29(9): 2034 – 2043.
- Incardona, J. P., Carls, M. G., Day, H. L., Sloan, C. A., Bolton, J. L., Collier, T. K., and Scholz, N. L. (2009). Cardiac arrhythmia is the primary response of embryonic Pacific herring (*Clupea pallasii*) exposed to crude oil during weathering. *Environmental Science and Technology*. 43(1): 201 – 207.
- Jones, E. (2019). Molecular Responses to Deepwater Horizon Oil Spill Contaminants in Estuarine Fish. PhD Dissertation, University of Southern Mississippi, 203pp
- Kadafa, A. A., Zakaria, M. P. and Othman, F. (2012). Oil spillage and pollution in Nigeria: organizational management and institutional framework. *Journal of Environment and Earth Science*. 2(4): 22 – 30.
- Kassahn, K. S., Caley, M. J., Ward, A. C., Connolly, A. R., Stone, G., and Crozier, R. H. (2007). Heterologous microarray experiments used to identify the early gene response to heat stress in a coral reef fish. *Molecular ecology*. 16(8): 1749 – 1763.
- Lee, R. F. and Anderson, J. W. (2005). Significance of cytochrome P450 system responses and levels of bile fluorescent aromatic compounds in marine wildlife following oil spills. *Mar Pollut Bull*. 50: 705 – 723.
- Lee, Y., Rhee, J., Hwang, D., Kim, I., Raisuddin, S. and Lee, J. (2008). p53 gene expression is modulated by endocrine-disrupting chemicals in the hermaphroditic fish, *Kryptolebias marmoratus*, *Comp. Physiol. Biochem. Toxicol. Pharmacol.* 147: 150 – 157.
- Liebermann, D. A. and Hoffman, B. (2008). Gadd45 in stress signaling. *Journal of molecular signaling*. 3(1): 1 – 8.
- Liu, M., Tee, C., Zeng, F., Sherry, J. P., Dixon, B., Bols, N. C. and Duncker, B. P. (2011). Characterization of p53 expression in rainbow trout. *Comp. Biochem. Physiol. C Toxicol. Pharmacol.* 154: 326 – 332.
- Min, J., Lee, S. K., and Gu, M. B. (2003). Effects of endocrine disrupting chemicals on distinct expression patterns of estrogen receptor, cytochrome p450 aromatase and p53 genes in *Oryzias latipes* liver. *J. Biochem. Mol. Toxicol.* 17: 272 – 277.
- Nebert, D. W., Dalton, T. P., Okey, A. B. and Gonzalez, F. J. (2004). Role of aryl hydrocarbon receptor-mediated induction of the CYP1 enzymes in environmental toxicity and cancer. *J. Biol. Chem.* 279: 23847–23850.
- Odiete, W.O. (2003). Standardization of test animals for toxicity evaluation for environmental regulation in Nigeria. *J. Nig. Environ. Soc.* 1: 340 – 350
- Olaifa, F. E. (2012). Bioassay using the water-soluble fraction of a Nigerian Light Crude oil on *Clarias gariepinus* fingerlings. *Niger. J. Physiol. Sci.* 27: 181–187
- Pfaffl, M. W., Horgan, G. H. and Dempfle L. (2002). Relative expression software tool (REST) for group-wide comparison and statistical analysis of relative expression results in real-time PCR. *Nucl. Acid Res.* 30 (9). Page e36, <https://doi.org/10.1093/nar/30.9.e36>
- Piña, B., Casado, M. and Quirós, L. (2007). Analysis of gene expression as a new tool in ecotoxicology and environmental monitoring. *TrAC Trends in Analytical Chemistry*. 26(11): 1145 – 1154.
- Raimondo, S., Montague, B. J. and Barron, M. G. (2007). Determinants of variability in acute to chronic toxicity ratios for aquatic invertebrates and fish. *Environmental Toxicology and Chemistry: An International Journal*. 26(9): 2019-2023.
- Rau-Embry, M., Billiard, S. M. and Di-Giulio, R. T. (2006). Lack of p53 induction in fish cells by model chemotherapeutics. *Oncogene*. 25: 2004–2010
- Sakuma, M. (1998). Probit analysis of preference data. *Applied entomology and zoology*, 33(3): 339 – 347.
- Santos, C., de Oliveira, M. T., de Syllos Cólus, I. M., Sofia, S. H., and dos Reis Martinez, C. B. (2018). Expression of *cyp1a* induced by benzo (A) pyrene and related biochemical and genotoxic biomarkers in the neotropical freshwater fish *Prochilodus lineatus*. *Environmental Toxicology and Pharmacology*. 61: 30 – 37.
- Santos, R. M., Weber, L., Souza, V. L., Soares, A. R. and Petry, A. C. (2016). Effects of water-soluble fraction of petroleum on growth and prey consumption of juvenile *Hoplias aff. Malabaricus* (Osteichthyes: Erythrinidae). *Braz. J. Biol.* 76 (1): 10 – 17.
- Sørhus, E., Donald, C. E., Da Silva, D., Thorsen, A., Karlsen, Ø. and Meier, S. (2021). Untangling

mechanisms of crude oil toxicity: linking gene expression, morphology and PAHs at two developmental stages in a cold-water fish. *Science of the Total Environment*. 757, 143896.

Sørhus, E., Incardona, J. P., Furmanek, T., Goetz, G. W., Scholz, N. L., Meier, S. and Jentoft, S. (2017). Novel adverse outcome pathways revealed by chemical genetics in a developing marine fish. *elife*, 6, e20707. <https://doi.org/10.7554/eLife.20707.001>.

Speidel, D. (2015). The role of DNA damage responses in p53 biology. *Archives of toxicology*. 89(4): 501-517.

Uno, T., Ishizuka, M., and Itakura, T. (2012). Cytochrome P450 (CYP) in fish. *Environmental toxicology and pharmacology*. 34(1): 1 – 13.

Vogelstein, B., Lane, D. and Levine, A. J. (2000). Surfing the p53 network. *Nature*. 408: 307 – 310.

Wedemeyer, G. A. (1997). Effects of rearing conditions on the health and physiological quality of fish

, A.D., Sumpter, J.P., Schreck, C.B. (Eds.), *Fish Stress and Health in Aquaculture*. Cambridge. University Press, Cambridge, pp. 35 – 72.

Whitehead, A., Dubansky, B., Bodinier, C., Garcia, T. I., Miles, S., Pilley, C. and Galvez, F. (2012). Genomic and physiological footprint of the Deepwater Horizon oil spill on resident marsh fishes. *Proceedings of the National Academy of Sciences*. 109(50): 20298 – 20302.

Williams, R. and Hubberstey, A. V. (2014). Benzo(a)pyrene exposure causes adaptive changes in p53 and CYP1A gene expression in Brown bullhead (*Ameiurus nebulosus*). *Aquat. Toxicol.* 156: 201 – 210.

Wincent, E., Jonsson, M. E., Bottai, M., Lundstedt, S. and Dreij, K. (2015). Aryl hydrocarbon receptor activation and developmental toxicity in zebrafish in response to soil extracts containing unsubstituted and oxygenated PAHs. *Environ. Sci. Technol.* 49: 3869 – 3877.




Figure 1: mRNA relative expression of *Cyp1a* and *p53* in the tissue of *C. gariepinus* after 90 days of exposure to WSFs of crude oil.

Table 1. Gene-specific primers used for RT PCR analysis

Gene	Mechanisms	Forward primer sequence	Reverse primer sequence
<i>CYP1A</i>	Detoxifying	GAAACCCGGCTGACTTCATT	GCCTCTGCATGAACACGTTA
<i>P53</i>	Tumour suppressor	TATACTGGCCTGTACGCATT G	TCTTGATAGTGAATAGACGG TCTTC
<i>b-actin</i>	Endogenous control	TGATGAAATCGCCGCACT	CCACAATGGATGGGAAGACA

Table 2: Mean values of Nanodrop quantification

Sample group	Nucleic acid	A260 (Abs)	A280 (Abs)	260/280	260/230
Control	157.00 ng/ μ l	2.14	1.42	1.49	1.77
30% LC ₅₀	255.55 ng/ μ l	5.11	3.63	1.68	1.92
45% LC ₅₀	247.75 ng/ μ l	2.96	1.87	1.40	1.62
60% LC ₅₀	182.00 ng/ μ l	3.64	2.31	1.58	1.77
75% LC ₅₀	212.45 ng/ μ l	4.25	2.65	1.59	1.80